arXiv:math/0606198v1 [math.GT] 8 Jun 2006

Classifying links under fused isotopy

ANDREW FISH and EBRU KEYMAN

School of Computing, Mathematical and Information Sciences, University of Brighton, UK
Andrew.Fish@brighton.ac.uk

Department of Mathematics, Middle East Technical University, Ankara, Turkey
ekeyman@metu.edu.tr

Abstract

All knots are fused isotopic to the unknot using a process known as virtualization. We extend and adapt this process to show that, up to fused isotopy, classical links are classified by their linking numbers.

Keywords: Fused isotopy, linking numbers

1. Introduction

Classical braids and links have been generalized to the virtual category [8] – adding virtual crossings and extending isotopy to allow the virtual analogues of the classical Reidemeister moves. The forbidden moves F_o and F_u, shown in Figure 1, are not allowable under virtual isotopy. Extending virtual isotopy in the virtual braid group VB_n to allow the F_o move gives rise to the welded braid group WB_n, which has been shown to be isomorphic to PC_n, the group of automorphisms of the free group on n elements of permutation-conjugacy type [3]. Allowing both of the forbidden moves F_o and F_u gives rise to fused isotopy [8]. That is, two virtual links L_1 and L_2 are called *fused isotopic* if L_2 can be obtained from L_1 by a finite sequence of Reidemeister moves, virtual moves and F_o, F_u moves.



Figure 1: The forbidden moves

Let $\mathcal{K}$ be the space of classical links embedded in S^3 and let $\mathcal{VK}$ be the space of virtual links. Kauffman [8], and independently Goussarov-Polyak-Viro [5], have shown that $\mathcal{K}$ embeds into $\mathcal{VK}$. Let f denote the natural inclusion of $\mathcal{VK}$ into the space of fused links $\mathcal{FK}$. Then we have $\mathcal{K} \overset{i}{\hookrightarrow} \mathcal{VK} \xrightarrow{f} \mathcal{FK}$, and when we refer to a classical link (under fused isotopy) we mean $f \circ i(L)$, the image of a link $L \in \mathcal{K}$ in the space $\mathcal{FK}$.

In [7], Kanenobu showed that all knots are fused isotopic to the unknot. He showed that all of the classical crossings of a virtual knot can be *virtualized*; that is every classical crossing can be changed into a virtual crossing by applying a sequence of fused isotopy moves. However, crossings between different components of a link cannot be virtualized using the same methods. The following theorem from [7] provides us with allowable moves under fused isotopy which were used in the virtualization procedure.

Theorem 1. The moves M_1, M_2 and M_3, shown in Figure 2, can be realised by fused isotopy. ☐



Figure 2: Allowable moves in fused isotopy

In [4] the authors showed that the Jones polynomial for welded and fused links is well-defined in a quotient of $\mathbb{Z}[A, A^{-1}]$ and observed that this polynomial depends only upon the linking number for links with two components. Inspired by this, we show that classical links, under fused isotopy, can be determined by the linking number of their components.

Theorem 2. A classical link L with n–components is completely determined by the linking numbers of each pair of components under fused isotopy.

The strategy that we use to prove Theorem 2 is to write L as the closure of a braid α on m strands (where $m \geq n$) and then to transform α into a pure braid β on n strands whose closure is also L. We show that β depends only on the linking numbers of the components of L. This means that any classical link with the same linking numbers as L can be obtained as the closure of β. We need some preliminaries before we proceed with the proof.

2. Preliminaries

Recall that an element of the pure braid group P_n is an n–strand braid where the permutation induced by the strings is the identity. P_n has a presentation with generators $A_{i,j}$ with $1 \leq i < j \leq n$ where

$$A_{i,j} = \sigma_{j-1}\sigma_{j-2}\ldots\sigma_i^2\ldots\sigma_{j-2}^{-1}\sigma_{j-1}^{-1} = \sigma_i^{-1}\sigma_{i+1}^{-1}\ldots\sigma_{j-1}^2\ldots\sigma_{i+1}\sigma_i.$$

Let U_k be the subgroup of P_n generated by $\{A_{i,k} \,:\, 1 \leq i < k\}$. Then every element of P_n can be written in the unique normal form $x_2x_3\ldots x_n$, where $x_k \in U_k$ (see [1] for details). Define $B_{i,j} := \sigma_{j-1}\ldots\sigma_{i+1}\sigma_i$ for $i < j$ and $B_{i,i} := 1$. Then by definition $A_{i,j+1} = B_{i,j}^{-1}A_{j,j+1}B_{i,j}$, and we can see from Figure 3 that for $k < i < j,\ B_{i,j}$ commutes with $A_{k,j+1}$ in B_n.



Figure 3: $A_{k,j+1}B_{i,j} = B_{i,j}A_{k,j+1}$

The virtual braid group on n–strands, VB_n, can be defined by adding extra generators τ_i, for $1 \leq i \leq n$, corresponding to the virtual crossings, and relations corresponding to the virtual isotopy moves [6, 8]. By adding the relations $\sigma_i^{-1}\tau_j\sigma_i = \sigma_j\tau_i\sigma_j^{-1}$ with $|i-j| = 1$, to the virtual braid group, we obtain the fused braid group, FB_n. If $j = i + 1$ the relation corresponds to the F_u move, and if $i = j + 1$ it corresponds to the F_o move. The explicit realization of the moves M_1, M_2 and M_3 using F_o and F_u moves is shown in [7], and this gives rise to the following consequences in FB_n:

$$\begin{array}{lrcl} M_1: & \sigma_i \tau_j \sigma_i & = & \sigma_j \tau_i \sigma_j \\ M_2: & \sigma_i^{-1} \tau_j \sigma_i^{-1} & = & \sigma_j^{-1} \tau_i \sigma_j^{-1} \\ M_3: & \sigma_i \sigma_j^{-1} \sigma_i & = & \sigma_j \sigma_i^{-1} \sigma_j, \end{array} \tag{2.1}$$

where $|i - j| = 1$.

The following lemmas are used in the proof of Theorem 2, and the indices have been chosen to match the usage in the proof. Let $\sim$ denote the equivalence class generated by fused isotopy and let $\widetilde{U}_k = U_k / \sim$.

Lemma 1. In FP_n we have:

$$A_{j,j+1} A_{i,j+1} A_{j,j+1}^{-1} = A_{i,j+1} \text{ where } 1 \leq i < j + 1 \leq n. \tag{2.2}$$

In other words, $A_{j,j+1} = \sigma_j^2$ is in the centre of $\widetilde{U}_{j+1}$.

Proof. Using the relations $\sigma_j \sigma_{j-1}^{-1} \sigma_j = \sigma_{j-1} \sigma_j^{-1} \sigma_{j-1}$ corresponding to an M_3 move, and $\sigma_j \sigma_{j-1} \sigma_j^{-1} = \sigma_{j-1}^{-1} \sigma_j \sigma_{j-1}$ corresponding to an R_3 move, we obtain

$$\begin{array}{rl} & \sigma_j^2 \sigma_{j-1}^{-1} \sigma_j^2 \sigma_{j-1} \sigma_j^{-2} \;=\; \sigma_j \left(\sigma_j \sigma_{j-1}^{-1} \sigma_j\right) \left(\sigma_j \sigma_{j-1} \sigma_j^{-1}\right) \sigma_j^{-1} \\ = & \sigma_j \left(\sigma_{j-1} \sigma_j^{-1} \sigma_{j-1}\right) \left(\sigma_{j-1}^{-1} \sigma_j \sigma_{j-1}\right) \sigma_j^{-1} \;=\; \sigma_j \sigma_{j-1}^2 \sigma_j^{-1}. \end{array} \tag{2.3}$$

Using Equation (2.3), we can compute

$$\begin{array}{rll} & A_{j,j+1} A_{i,j+1} A_{j,j+1}^{-1} & \\ = & \sigma_j^2 B_{i,j}^{-1} A_{j,j+1} B_{i,j} \sigma_j^{-2} & \\ = & B_{i,j-1}^{-1} \sigma_j^2 \sigma_{j-1}^{-1} \sigma_j^2 \sigma_{j-1} \sigma_j^{-2} B_{i,j-1} & \text{by commutation in } B_n \\ = & B_{i,j-1}^{-1} \sigma_j \sigma_{j-1}^2 \sigma_j^{-1} B_{i,j-1} & \text{by Equation (2.3)} \\ = & \sigma_j B_{i,j-1}^{-1} \sigma_{j-1}^2 B_{i,j-1} \sigma_j^{-1} & \text{by commutation in } B_n \\ = & \sigma_j A_{i,j} \sigma_j^{-1} & \\ = & A_{i,j+1}. & \square \end{array}$$

Lemma 2. For every $1 \leq j + 1 < n$, the subgroup $\widetilde{U}_{j+1}$ of FP_n is commutative.

Proof. Assume without loss of generality that $k < i$. Then

$$\begin{array}{rcll} A_{k,j+1} A_{i,j+1} & = & A_{k,j+1} B_{i,j}^{-1} A_{j,j+1} B_{i,j} & \\ & = & B_{i,j}^{-1} A_{k,j+1} A_{j,j+1} B_{i,j} & \text{by commutation in } B_n \\ & = & B_{i,j}^{-1} A_{j,j+1} A_{k,j+1} B_{i,j} & \text{by Lemma 1} \\ & = & B_{i,j}^{-1} A_{j,j+1} B_{i,j} A_{k,j+1} & \text{by commutation in } B_n \\ & = & A_{i,j+1} A_{k,j+1}. & \square \end{array}$$

Lemma 3. In FB_n we have:

$$A_{i,j+1} \tau_j = \tau_j A_{i,j} \qquad \text{for } 1 \leq i \leq j - 1. \tag{2.4}$$

Proof. Using the relations $\sigma_j\sigma_{j-1}\tau_j = \tau_{j-1}\sigma_j\sigma_{j-1}$ corresponding to an F_0 move, and $\sigma_{j-1}^{-1}\sigma_j\tau_{j-1} = \tau_j\sigma_{j-1}\sigma_j^{-1}$ corresponding to an M_1 move, we obtain

$$\begin{aligned}
A_{i,j+1}\tau_j &= B_{i,j}^{-1}\sigma_j^2 B_{i,j}\tau_j \\
&= B_{i,j-1}^{-1}\sigma_{j-1}^{-1}\sigma_j\sigma_j\sigma_{j-1}B_{i,j-1}\tau_j \\
&= B_{i,j-1}^{-1}\sigma_{j-1}^{-1}\sigma_j\sigma_j\sigma_{j-1}\tau_j B_{i,j-1} && \text{by commutation in } VB_n \\
&= B_{i,j-1}^{-1}\sigma_{j-1}^{-1}\sigma_j\tau_{j-1}\sigma_j\sigma_{j-1}B_{i,j-1} && \text{by an } F_o \text{ move} \\
&= B_{i,j-1}^{-1}\tau_j\sigma_{j-1}\sigma_j^{-1}\sigma_j\sigma_{j-1}B_{i,j-1} && \text{by an } M_1 \text{ move} \\
&= \tau_j B_{i,j-1}^{-1}\sigma_{j-1}^2 B_{i,j-1} && \text{by commutation in } VB_n \\
&= \tau_j A_{i,j}. && \square
\end{aligned}$$

3. Proof of Theorem 2

Let L be a classical link with n–components. By Alexander's Theorem, there exists $\alpha \in B_m$ with $m \geq n$, such that the closure of α is L. Chow [2] (or see page 22 of [1]) shows that every α can be written in the form $\alpha = x_2 B_{k_2,2}\dots x_m B_{k_m,m}$ where $x_i \in U_i \leq P_m$ and $1 \leq k_i \leq i$. Let $\hat{\alpha}$ denote the closure of α. If $m > n$ then we will construct $\beta \in B_n$ such that $\hat{\beta}$ is fused isotopic to $\hat{\alpha}$.

If $B_{k_i,i} = 1$ for all $i = 2,\dots,m$ then α is a pure braid, which means that m must be equal to n. So let us assume that $B_{k_s,s} \neq 1$, for some s, and that if $i > s$ then $B_{k_i,i} = 1$. This means that the permutation induced by α is the identity on the strands $s+1,\dots,m$. Therefore, each of these strands forms a separate component of the link $\hat{\alpha}$. Now, conjugating α with $B_{1,m}$ gives $B_{1,m}^{-1}\alpha B_{1,m}$, and as shown in Figure 4, the $(m-1)$–strand of the original braid α becomes the m–strand of the new braid.



Figure 4: $B_{1,m}^{-1}\alpha B_{1,m}$

Thus if we conjugate α with $B_{1,m}$ $(m-s)$ times, we get $\alpha' = B_{1,m}^{s-m}\alpha B_{1,m}^{m-s}$ and the s–strand of α becomes the m–strand of α'. Since α' is just a conjugate of α their closures are isotopic. Now write $\alpha' = y_2 B_{t_2,2}\dots y_m B_{t_m,m}$ with $y_i \in \widetilde{U}_i$. Then $B_{t_m,m} \neq 1$ and so by definition, $B_{t_m,m} = \sigma_{m-1}B_{t_m,m-1}$. A picture of α' is shown in Figure 5, where $W = y_2 B_{t_2,2}\dots y_{m-1}B_{t_{m-1},m-1}$.

Since $\widetilde{U}_m$ is commutative (by Lemma 2), we can write

$$y_m = \underbrace{A_{1,m}^{r_1}\dots A_{m-2,m}^{r_{m-2}}}_{v_m}\underbrace{A_{m-1,m}^{r_{m-1}}}_{\sigma_{m-1}^{2r_{m-1}}}$$

for some $r_1,\dots,r_{m-1}$. By definition, $A_{m-1,m}^{r_{m-1}} = \sigma_{m-1}^{2r_{m-1}}$, and since $B_{t_m,m} = \sigma_{m-1}B_{t_m,m-1}$, we obtain $y_m B_{t_m,m} = v_m\sigma_{m-1}^{2r_{m-1}+1}B_{t_m,m-1}$, where $v_m = A_{1,m}^{r_1}\dots A_{m-2,m}^{r_{m-2}}$.



Figure 5: The braid α'

Since W does not involve the m–strand and y_m is a pure braid, Figure 5 shows that the m–strand and the other strand that is involved in the last occurrence (and hence in all of the previous occurrences) of σ_{m-1} in α' belong to the same component of $L = \hat{\alpha}'$. Therefore, following the strategy in [7], we can virtualize all of the $2r_{m-1}+1$ crossings in $\hat{\alpha}'$ which correspond to $\sigma_{m-1}^{2r_{m-1}+1}$ in α'. In doing so we have not changed the fused isotopy class of L but we have obtained L as the closure of $\alpha_1 = Wu_m\tau_{m-1}^{2r_{m-1}+1}B_{t_m,m-1} = Wu_m\tau_{m-1}B_{t_m,m-1}$. By Lemma 3, we obtain

$$\alpha_1 = W\tau_{m-1}v_{m-1}B_{t_m,m-1}$$

where $v_{m-1} = A_{1,m-1}^{r_1} \dots A_{m-2,m-1}^{r_{m-2}}$.

Figure 6 shows that there is only one crossing involving the m–strand in the braid α_1. This is the occurrence of τ_{m-1}. In $\hat{\alpha}_1$, we can get rid of the virtual crossing corresponding to τ_{m-1} with a virtual move (of type I). We have obtained a new link diagram $\hat{\alpha}_2$ where $\alpha_2 = Wv_{m-1}B_{t_m,m-1}$ has $m-1$ strands and $\hat{\alpha}_2$ is fused isotopic to L.



Figure 6: The braid α_1

If we continue with this process, eventually we will get a braid β in B_n whose closure is fused isotopic to L. Note that each strand of β corresponds to a different component of L and therefore β must be a pure braid. For $i < j$, define the group homomorphism $\delta_{i,j} : PB_n \to \mathbb{Z}$ by

$$\delta_{i,j}(A_{s,t}) = \begin{cases} 1 & \text{if } s = i \text{ and } t = j \\ 0 & \text{otherwise.} \end{cases}$$

Since β is a pure braid it is easy to see that $\delta_{i,j}(\beta) = lk(\ell_i, \ell_j)$ where ℓ_i and ℓ_j are the corresponding components of $\hat{\beta}$.

This proves that any classical link L with n–components can be obtained as the closure of a pure braid $\beta = x_2 \dots x_n$ where each x_k can be written in the form $x_k = A_{1,k}^{\delta_{1,k}} \dots A_{k-1,k}^{\delta_{k-1,k}}$. This shows that β depends only on the linking number of the components. □

Remark 1. We do not believe that the theorem will generalize to non-classical links where there are virtual crossings between different components. For example, let U_2 be the trivial link with two components and let $L = \hat{\alpha}$ where $\alpha = \sigma_1\tau_1\sigma_1^{-1}\tau_1 \in FB_2$. Then both of these links have linking number 0 but we conjecture that they are not fused isotopic (although currently there are no known invariants to distinguish them).

[1] J. Birman, *Braids, links and mapping class groups*, Annals of Mathematical Studies **82**, Princeton Univ. Press (1974).

[2] W.L. Chow, *On the algebraic braid group*. Annals of Math. **49** (1948), 654–658.

[3] R. Fenn, R. Rimányi and C. Rourke, *The braid-permutation group*, Topology **36** (1997), 123–135.

[4] A. Fish and E. Keyman, *Jones polynomial invariants*, JKTR **15:3** (2006) 339–350.

[5] M. Goussarov, M. Polyak and O. Viro, *Finite-type invariants of classical and virtual knots*, Topology **39** (2000) 1045–1068.

[6] S. Kamada, *Braid presentation of virtual knots and welded knots*, arXiv:math.GT/0008092 v1.

[7] T. Kanenobu, *Forbidden moves unknot a virtual knot*, JKTR **10:1** (2001) 89–96.

[8] L.H. Kauffman, *Virtual Knot Theory*, European J. Combinatorics **20** (1999) 663–690.